## **Aftercare Information for a One-Day Abortion Procedure**

The following information is for intended for patients of Hope Clinic following a one-day in-clinic abortion procedure.

Please reference this information as needed throughout your recovery. (Version 1.2023)

hopeclinic.com | (618) 451-5722 | info@hopeclinic.com

## What to Expect

#### Vaginal Bleeding:

- Bleeding is expected to start and stop randomly for 2-4 weeks after an abortion procedure.
- The color of the blood can range from red to brown.
- We recommend wearing heavy protection or overnight menstrual pads for at least 48 hours after your procedure.
- As long as bleeding is not exceeding 2 menstrual pads in 1 hour, your bleeding is considered normal.
- For most people, a menstrual period returns in 4-6 weeks after an abortion.

#### Cramps/Pain:

- Can vary from mild to moderate; usually lasts for about 1 week after your procedure.
- Take pain medications as directed. Other ways you can relieve pain include using a heating pad, taking a shower, taking a hot bath, having someone rub your back, in general, you can do whatever you would normally do for period cramps.

### What to Avoid

#### **Sex & Tampons:**

- It is up to you to determine when you feel comfortable having sex again. Please avoid having sex until you feel ready to do so.
- Avoid using tampons for 48 hours after your procedure.
  - o After 48 hours, you can use whatever products you feel most comfortable using.

#### **Activities:**

- Most people can return to their regular level of physical activity one day after their procedure. However, reduce any activity that increases your cramping and/or bleeding.
- If you received IV sedation, it is important that you do NOT drive for at least 24 hours after your procedure.

## **Medications**

#### Pain Management:

• **Ibuprofen (Advil or Motrin) 600mg:** every 6 hours as needed for pain/cramping

#### Bleeding:

- Some patients will be discharged with additional medication to prevent bleeding, as needed.
  - Misoprostol (200mcg): one tablet under tongue every
     3 hours until pills are gone.

#### **Antibiotics:**

 You received a 1-time preventative dose of antibiotic at your appointment. Unless directed otherwise by your provider, you do not need additional antibiotics.

#### **Birth Control:**

 You may start your preferred contraceptive method immediately. More information is available on our website.

## WARNING SIGNS

The following could be signs of problems which could lead to severe health issues such as life-threatening hemorrhage or infection.

Please call the emergency number *immediately* if you have any of the following:

- Heavy bleeding: soaking through (from top to bottom/end to end)
   2 super maxi pads in an hour for 2 or more hours
- Fever: a temperature of 100.4 degrees or greater
- Severe pain: with NO relief from pain medications
- Nausea/vomiting/diarrhea/dizziness/extreme fatigue or weakness

# Emergency Contact Number (618) 346-3040

The Hope Clinic doctors/nurses are available 24 hours a day, every day, for emergency calls.

## Follow-Up

A follow-up is an optional check-up after your procedure. If you feel something is not quite right, or even if you just want peace of mind, you can call to schedule a follow-up at Hope Clinic or with another healthcare provider. You can follow up even if you are still bleeding.